

Instructions for Parents
For Care Following

Ureteroneocystostomy

ACTIVITY

Your child should gradually increase his/her activity over the next 1-2 weeks. Your child should not participate in organized sports or gym class for 3 weeks.

DIET

You may feed your child whatever he/she wants to eat.

DRESSINGS

The incision is closed with dissolving stitches. Small strips of tape called SteriStrips cover the incision line. These tapes will fall off within 2 weeks after the surgery. Sometimes a **stent** (small catheter) is left in place coming out of the skin of the flank. If a stent was left, clean around the point where the tube comes through the skin and around the blue plastic plate around the stent twice daily with hydrogen peroxide. You may use a swab for this. Change the gauze over the incision twice daily.

MEDICATION

Your child will need pain medication following the surgery. Unless you were instructed otherwise, you can use ibuprofen (Motrin for children, Advil for children, or a generic form). You can give your child _____ mg. Ibuprofen every 6 hours. If ibuprofen does not relieve the pain well enough, use the Tylenol with codeine according to the directions on the bottle. Use either ibuprofen or Tylenol with codeine, not both. You should continue to give your child antibiotics until you return to the clinic for follow-up. Follow the directions on the bottle of antibiotics.

BATHING

If no stent was left following the surgery, your child may bathe or shower 48 hours following the surgery. For the first 7 days following the surgery limit the time in the tub or shower to 5 – 10 minutes. You may let the water run over the incision. If a stent was left at surgery, your child should have a sponge bath until the stent/drain is removed. The stent is usually removed 1 – 2 weeks following the surgery. Be sure to give your child a dose of pain medication just before coming to the clinic for removal of the stent.

ADDITIONAL INSTRUCTIONS

If the incision becomes red, swollen or hot, or if blood, clear or creamy fluid comes out of the incision or through the point where the stent or drain comes through the skin, please call our office at the number below. Your child may have a low-grade fever the night of surgery. Call your doctor if the fever continues or if it is higher than 102 degrees Fahrenheit. If your child develops nausea or is unable to eat, please call our office.

If your child should have difficulty with breathing, bleeding that you feel is excessive, persistent nausea or vomiting, pain that is unusual, swelling, or fever, please call your surgeon at 708/216-6266 (Monday-Friday 8:30 a.m. to 5:00 PM) or 708/216-8000 (evenings, nights or weekends). **In the event of an urgent emergency, you may take your child to the Loyola University Medical Center Emergency Department (708/216-3882) or to an emergency room near to you.**

Patient/Parent/Guardian

Nurse/Physician

Date

David A. Hatch, M.D., F.A.C.S., F.A.A.P.

Derek Matoka, M.D.

Section of Pediatric Urology
Loyola University Medical Center
2160 S. First Avenue
Maywood, IL 60153

Anne Casaccio, RN 708/216-5111

Linda Bauman, Secretary 708/216-6266

lbauman@luhs.org

Office:

708/216-6266

Appointments:

Central Scheduling (all clinics) 708/216-8563

Darien 630/985-4989

Elmhurst 708/327-7030


Hickory Hills 708/233-5333

Homer Glen 708.645.3400

Oakbrook Terrace 630/953-6600

Wheaton 603.665.5995

Directions and maps available at www.luhs.org



Instructions for Parents
For Care Following

Ureteroneocystostomy

ACTIVITY

Your child should gradually increase his/her activity over the next 1-2 weeks. Your child should not participate in organized sports or gym class for 3 weeks.

DIET

You may feed your child whatever he/she wants to eat.

DRESSINGS

The incision is closed with dissolving stitches. Small strips of tape called SteriStrips cover the incision line. These tapes will fall off within 2 weeks after the surgery. Sometimes a stent (small catheter) is left in place coming out of the skin of the flank. If a stent was left, clean around the point where the tube comes through the skin and around the blue plastic plate around the stent twice daily with hydrogen peroxide. You may use a swab for this. Change the gauze over the incision twice daily.

MEDICATION

Your child will need pain medication following the surgery. Unless you were instructed otherwise, you can use ibuprofen (Motrin for children, Advil for children, or a generic form). You can give your child _____ mg. Ibuprofen every 6 hours. If ibuprofen does not relieve the pain well enough, use the Tylenol with codeine according to the directions on the bottle. Use either ibuprofen or Tylenol with codeine, not both. You should continue to give your child antibiotics until you return to the clinic for follow-up. Follow the directions on the bottle of antibiotics.

BATHING

If no stent was left following the surgery, your child may bathe or shower 48 hours following the surgery. For the first 7 days following the surgery limit the time in the tub or shower to 5 – 10 minutes. You may let the water run over the incision. If a stent was left at surgery, your child should have a sponge bath until the stent/drain is removed. The stent is usually removed 1 – 2 weeks following the surgery. Be sure to give your child a dose of pain medication just before coming to the clinic for removal of the stent.

ADDITIONAL INSTRUCTIONS

If the incision becomes red, swollen or hot, or if blood, clear or creamy fluid comes out of the incision or through the point where the stent or drain comes through the skin, please call our office at the number below. Your child may have a low-grade fever the night of surgery. Call your doctor if the fever continues or if it is higher than 102 degrees Fahrenheit. If your child develops nausea or is unable to eat, please call our office.

If your child should have difficulty with breathing, bleeding that you feel is excessive, persistent nausea or vomiting, pain that is unusual, swelling, or fever, please call your surgeon at 708/216-6266 (Monday-Friday 8:30 a.m. to 5:00 PM) or 708/216-8000 (evenings, nights or weekends). **In the event of an urgent emergency, you may take your child to the Loyola University Medical Center Emergency Department (708/216-3882) or to an emergency room near to you.**

Patient/Parent/Guardian

Nurse/Physician

Date

David A. Hatch, M.D., F.A.C.S., F.A.A.P.
Derek Matoka, M.D.
Section of Pediatric Urology
Loyola University Medical Center
2160 S. First Avenue
Maywood, IL 60153

Anne Casaccio, RN 708/216-5111
Linda Bauman, Secretary 708/216-6266
lbauman@luhs.org

Office: 708/216-6266
Appointments:
Central Scheduling (all clinics) 708/216-8563
Darien 630/985-4989
Elmhurst 708/327-7030
Hickory Hills 708/233-5333
Homer Glen 708.645.3400
Oakbrook Terrace 630/953-6600
Wheaton 603.665.5995

Directions and maps available at www.luhs.org