

CLINICAL PRACTICE

## Palliative Care

R. Sean Morrison, M.D., and Diane E. Meier, M.D.

From the Hertzberg Palliative Care Institute, Brookdale Department of Geriatrics and Adult Development, Mount Sinai School of Medicine, New York. Address reprint requests to Dr. Morrison at the Department of Geriatrics, Box 1070, Mount Sinai School of Medicine, One Gustave L. Levy Place, New York, NY 10029, or at sean.morrison@mssm.edu.

N Engl J Med 2004;350:2582-90.  
Copyright © 2004 Massachusetts Medical Society.

**An 85-year-old man with New York Heart Association class IV heart failure, hypertension, and moderate Alzheimer's disease is admitted to the hospital after a hip fracture. His postoperative course is complicated by pneumonia, delirium, and pressure ulcers on his heels and sacrum. He is losing weight and is unable to participate in rehabilitation because of his confusion. This is his fourth hospitalization in the past year. His 84-year-old wife, who has been caring for him at home, feels overwhelmed by his medical and personal care needs. The patient's physician is increasingly frustrated by his frequent readmissions. What might she do to address his needs, alleviate his suffering, and facilitate his discharge from the hospital and subsequent care at home?**

### THE CLINICAL PROBLEM

By 2030, 20 percent of the U.S. population will be over the age of 65 years.<sup>1</sup> For most people, the years after the age of 65 are a time of good health, independence, and integration of a life's work and experience. Eventually, most adults will have one or more chronic illnesses with which they will live for years before they die. These years are often characterized by physical and psychological distress, progressive functional dependence and frailty, and increased needs for family and external support.<sup>1</sup> Studies suggest that medical care for patients with serious and advanced illnesses is characterized by the undertreatment of symptoms, conflict about who should make decisions about the patient's care, impairments in caregivers' physical and psychological health, and depletion of family resources.<sup>2-5</sup>

### STRATEGIES AND EVIDENCE

#### THE ROLE OF PALLIATIVE CARE

There are many reasons why patients who have advanced illnesses receive inadequate care, but most of those reasons are rooted in a medical philosophy that is focused almost exclusively on curing illness and prolonging life, rather than on improving the quality of life and relieving suffering. Traditionally, medical care has been articulated as having two mutually exclusive goals: either to cure disease and prolong life or to provide comfort care.<sup>1</sup> Given this dichotomy, the decision to focus on reducing suffering is made only after life-prolonging treatment has been ineffectual and death is imminent.<sup>6</sup> In the United States, this forced choice is driven largely by the reimbursement system — that is, regular Medicare covers curative therapies and the Medicare hospice benefit covers comfort care. That division of services results both in the provision of burdensome and costly life-prolonging treatments when they are no longer beneficial and in preventable suffering during all stages of advanced illness.<sup>7</sup> In contrast, patients would benefit most from care that included a combination of life-prolonging treatment (when possible and appropriate), palliation of symptoms, rehabilitation, and support for caregivers.

Meeting the needs of patients will require that physicians employ skills that are


not traditionally taught in medical schools.<sup>8</sup> Palliative care aims to relieve suffering and improve the quality of life for patients with advanced illnesses and their families through specific knowledge and skills, including communication with patients and family members; management of pain and other symptoms; psychosocial, spiritual, and bereavement support; and coordination of an array of medical and social services.<sup>1</sup> Palliative care should be offered simultaneously with all other medical treatment.

**PHYSICIAN-PATIENT COMMUNICATION**

Communicating with patients is a core skill of palliative medicine.<sup>9</sup> Studies suggest that in a typical clinical encounter, clinicians elicit fewer than half of patients' concerns and consistently fail to discuss patients' values, goals of care, and preferences with regard to treatment.<sup>10</sup> Empirical evidence supports the effectiveness of clinicians' use of specific communication skills in enhancing disclosure of the issues of concern to a patient, decreasing anxiety, assessing depression, and improving a patient's well-being and the level of the patient's and the family's satisfaction with the treatment.<sup>10</sup> Those communication skills include making eye contact with patients, asking open-ended questions, responding to a patient's affect, and demonstrating empathy. When a physician is informing a patient about a poor prognosis, conducting an open, "patient-centered" interview (in which the emphasis is on empathy, openness, and reassurance), rather than the traditional closed, "physician-centered" interview (in which the physician is focused on the task), has been associated with an improved level of satisfaction on the part of patients and their families.<sup>11</sup>

Guidelines have been developed for establishing goals of medical care, communicating bad news, and withholding or withdrawing medical treatments (Fig. 1).<sup>12,13</sup> Whether the use of these protocols actually helps patients and their families has not been empirically determined, but we and others<sup>12-16</sup> have found these guidelines to be useful in clinical practice. One randomized trial in the United Kingdom showed that a postgraduate course in communication skills that had cognitive, behavioral, and affective components led to improved communication skills among the physicians.<sup>17</sup> Similar courses are offered annually in the United States.<sup>18,19</sup>

Palliative care begins with establishing the goals



of care. Outlining realistic and attainable goals assumes an increased importance in the setting of advanced disease, in which treatments intended to cure the disease and prolong life may be more burdensome than beneficial. Whereas the goal for

some patients may be to prolong life at any cost, studies suggest that what most seriously ill patients want is to have their pain and other symptoms relieved, improve their quality of life, avoid being a burden to their family, have a closer relationship with loved ones, and maintain a sense of control.<sup>9,20</sup>

Establishing clear goals can facilitate decision making regarding treatment. For example, in the case of a patient with end-stage dementia who has dysphagia, placement of a percutaneous endoscopic gastrostomy (PEG) tube may be considered. If the primary goal is to reduce suffering and enhance the quality of life, then placing a PEG tube is unlikely to meet these goals, since it requires a painful, invasive procedure; eliminates the pleasurable oral sensations of eating and drinking; is associated with an increased use of restraints; can cause cellulitis, vomiting, diarrhea, and fluid imbalances; and is unlikely to reduce the risk of aspiration.<sup>21</sup> Even in cases in which a primary goal is to prolong life, the role of a PEG tube remains questionable; no survival benefits have been shown in observational studies in which patients with dementia who received feeding tubes were compared with similar patients who did not receive feeding tubes.<sup>21</sup>

Expert opinion suggests that clinicians can assist patients and their families in establishing their own goals by means of open-ended and probing questions. Some examples of the types of questions include “What makes life worth living for you?” “Given the severity of your illness, what are the most important things for you to achieve?” “What are your most important hopes?” “What are your biggest fears?” and “What would you consider to be a fate worse than death?”<sup>16</sup> The goals that patients establish may overlap, be contradictory, rise and fall in importance, and shift with the progression of the disease.<sup>12,16</sup> Warning signs of poorly established or conflicting goals can include frequent and lengthy hospitalizations; feelings of frustration, anger, or powerlessness on the part of the physician; and feelings of being burdened on the part of caregivers.<sup>15</sup>

Once goals are established, they can be used to construct advance directives about the types of care that patients want. Most studies,<sup>22</sup> although not all,<sup>23,24</sup> have shown that few patients have advance directives and that the documents they do have are relatively ineffectual in enhancing physician–patient communication, facilitating decision making about resuscitation,<sup>25</sup> or influencing terminal care in hospitals.<sup>22</sup> It is possible that the gradual rise in the

prevalence of advance directives over the past decade will improve their effectiveness, as physicians and patients become more familiar with them and physicians become more comfortable using them for assistance in guiding the care of cognitively impaired adults. A recent report suggested that the focus of advance care planning should shift from discussing specific treatments to defining an acceptable quality of life and setting goals for care under various likely clinical scenarios.<sup>26,27</sup> Whether this goal-centered approach to advance care planning will affect clinical outcomes is unknown.

#### ASSESSMENT AND TREATMENT OF SYMPTOMS

A fundamental goal of palliative care is the relief of pain and other symptoms.<sup>9,20</sup> Successful approaches to the assessment and management of pain and some physical and psychological symptoms have been established in controlled trials.<sup>12,28–30</sup> Despite these advances, undertreatment of symptoms persists in the majority of patients and settings.<sup>2,31</sup>

Relief of suffering begins with routine and standardized symptom assessment with use of validated instruments. Routine assessment has been shown to identify overlooked and unreported symptoms, facilitate treatment, and enhance patient and family satisfaction.<sup>32,33</sup> Clinically useful assessment instruments can be found on the Web sites of the Center to Advance Palliative Care ([www.capc.org](http://www.capc.org)) and of Brown University’s Center for Gerontology and Health Care Research, which features a tool kit of instruments to measure end-of-life care ([www.chcr.brown.edu/pcoc/toolkit.htm](http://www.chcr.brown.edu/pcoc/toolkit.htm)).

Improved treatment of symptoms has been associated with the enhancement of patient and family satisfaction, functional status, quality of life, and other clinical outcomes.<sup>32,34</sup> Although a comprehensive review of strategies for assessment and therapy of symptoms is beyond the scope of this review, Table 1 summarizes approaches to commonly encountered symptoms in advanced illnesses.

#### PSYCHOSOCIAL, SPIRITUAL, AND BEREAVEMENT SUPPORT

Providing psychosocial, spiritual, and bereavement support to patients and caregivers is a key component of palliative care. Patients who experience spiritual and psychological distress are more likely to express a desire for death than are other patients,<sup>42</sup> and their family members are more likely to have an extended or complicated grief and bereavement process (defined as grief that lasts for

**Table 1. Approaches to the Management of Pain and Other Common Symptoms.\***

Symptom	Assessment	Treatment
<b>Anorexia and cachexia</b>	Is a disease process causing the symptom, or is it secondary to other symptoms (e.g., nausea and constipation) that can be treated? Is the patient troubled by the symptom?	Consider megestrol acetate or dexamethasone.
<b>Anxiety</b>	Does the patient exhibit restlessness, agitation, insomnia, hyperventilation, tachycardia, or excessive worry?	Recommend supportive counseling and consider prescribing benzodiazepines (in the elderly, avoid benzodiazepines with long half-lives).
<b>Constipation</b>	Is the patient taking opioids? Does the patient have a fecal impaction?	Prescribe a stool softener (ineffective alone) plus escalating doses of a stimulant; if escalation of the dose is ineffectual, agents from other classes (e.g., osmotic laxatives and enemas) should be added.
<b>Depression</b>	How does the patient respond to the question "Are you depressed?" Does the patient express or exhibit any of the following feelings: helplessness, hopelessness, anhedonia, loss of self-esteem, worthlessness, persistent dysphoria, and suicidal ideation? (Somatic symptoms are not reliable indicators of depression in this population.)	Recommend supportive psychotherapy, cognitive approaches, behavioral techniques, pharmacologic therapies, or a combination of these interventions; prescribe psychostimulants for rapid treatment of symptoms (within days) or selective serotonin-reuptake inhibitors, which may require three to four weeks to take effect; tricyclic antidepressants are relatively contraindicated because of their side effects.
<b>Delirium</b>	Was the onset of confusion acute? Is the patient disoriented or experiencing changes in the level of consciousness or minute-to-minute fluctuations? Is the condition reversible?	Identify underlying causes and manage symptoms; recommend behavioral therapies, including avoidance of excess stimulation, frequent reorientation, and reassurance; ensure presence of family caregivers; prescribe haloperidol, risperidone, or olanzapine. (Chlorpromazine can be used for agitated or terminal delirium, but benzodiazepines have been found to exacerbate delirium and should be avoided.)
<b>Dyspnea</b>	Does the symptom have reversible causes?	Prescribe oxygen to treat hypoxia-induced dyspnea or to provide symptomatic relief, when hypoxia is absent, through stimulation of the V2 branch of the trigeminal nerve. Opioids relieve breathlessness without measurable reductions in respiratory rate or oxygen saturation; effective doses are often lower than those used to treat pain. Consider anxiolytics (e.g., low-dose benzodiazepines) and use reassurance, relaxation, distraction, and massage therapy.
<b>Nausea</b>	Which mechanism is causing the symptom (e.g., stimulation of the chemoreceptor trigger zone, gastric stimulation, delayed gastric emptying or "squashed stomach" syndrome, bowel obstruction, intracranial processes, or vestibular vertigo)?	Prescribe an agent directed at the underlying cause. Multiple agents directed at various receptors or mechanisms may be required.
<b>Pain</b>	How severe is the symptom (as assessed with the use of validated instruments)?	Prescribe medications to be administered on a standing or regular basis; as-needed or rescue doses should be available for breakthrough pain or pain not controlled by the standing regimen; start a regimen to prevent constipation for all patients receiving opioids. For mild pain: use acetaminophen or a nonsteroidal antiinflammatory agent (consider opioids in older adults). For moderate pain: titrate short-acting opioids. For severe pain: rapidly titrate short-acting opioids until pain is relieved or intolerable side effects develop; start long-acting opioids (e.g., sustained-release morphine or oxycodone and transdermal fentanyl) once pain is well controlled; use methadone only if experienced in its use. Rescue doses: prescribe immediate-release opioids consisting of 10% of the 24-hour total opioid dose to be given every hour (orally) or every 30 minutes (parenterally) as needed. Concomitant analgesics (e.g., corticosteroids, anticonvulsants, tricyclic antidepressants, and bisphosphonates) should be used when applicable.

\* For details of recommendations and more information about the management of symptoms, see Foley,<sup>35</sup> American Geriatrics Society Panel on Persistent Pain in Older Persons,<sup>36</sup> Luce and Luce,<sup>37</sup> Casarett and Inouye,<sup>38</sup> Strasser and Bruera,<sup>39</sup> Block,<sup>40</sup> and Regnard and Comiskey.<sup>41</sup>

at least 14 months after the death and results in a failure of the survivor to return to his or her normal activities) and are at higher risk for illness and death.<sup>43</sup>

Studies suggest that patients welcome inquiries about their spiritual well-being from their physicians,<sup>44</sup> although interventions to address spiritual distress have not been well developed or well evaluated. Support groups have been found to reduce stress and depression experienced by caregivers.<sup>45,46</sup> Although many formal interventions to address complicated grief and bereavement have been described, data are lacking on their effects on outcomes.<sup>47</sup> Two recent studies reported lower morbidity and mortality<sup>48</sup> and better emotional support<sup>49</sup> among surviving family members of hospice patients than among family members of patients who did not receive hospice services, although it is uncertain whether this difference reflects the nature of families who elect hospice care rather than the effects of the intervention.<sup>48</sup>

#### COORDINATION OF CARE

Several studies demonstrate that the personal and practical care needs of patients who are seriously ill and their families are not adequately addressed by routine office visits or hospital and nursing home stays and that this failure results in substantial burdens — medical, psychological, and financial — on patients and their caregivers.<sup>3-5,49,50</sup> In the context of chronic progressive disease, the ability of physicians to coordinate an array of social and medical services on behalf of patients and families assumes increased importance.

Various comprehensive care programs are available to help physicians manage the care of their patients who have serious and complex illnesses. Palliative care programs within home care organizations, hospitals, and nursing homes are increasingly prevalent in the United States and provide comprehensive interdisciplinary care for patients and families in collaboration and consultation with primary care physicians.<sup>51</sup> Hospice services, under the Medicare benefit, are available in most U.S. communities and provide palliative care, primarily at home, for patients with a life expectancy of six months or less who are willing to forgo insurance coverage for life-prolonging treatments. Studies suggest that referral to palliative care programs and hospice results in beneficial effects on patients' symptoms, reduced hospital costs, a greater likelihood of death at home, and a higher level of patient

and family satisfaction than does conventional care.<sup>49,50,52,53</sup>

Other programs that coordinate care for patients who have complex illnesses are available in most communities. Programs that coordinate home care services for patients with chronic conditions (e.g., depression, heart failure, or the acquired immunodeficiency syndrome) have been associated with increased patient and family satisfaction and reductions in mortality, use of hospital services, visits to physicians' offices, and admission to nursing homes.<sup>54-56</sup> Case management is usually provided by managed-care organizations, Medicare managed-care services, commercial insurers, some home health care agencies, and privately hired case managers. The quality, cost, and extent of the services provided are highly variable.

Comprehensive multidisciplinary home care programs that serve frail older adults have been evaluated under Medicare and the Veterans Health Administration (VHA). The Program of All-Inclusive Care for the Elderly (PACE) is a capitated Medicare and Medicaid benefit for frail older adults that offers comprehensive medical and social services at 25 adult day health centers, in homes, and at inpatient facilities. (More information about the program is available online at [www.cms.hhs.gov/pace/](http://www.cms.hhs.gov/pace/).) Patients who use the services provided by PACE have higher rates of completing advance directives for their care<sup>57</sup> and lower rates of admission to nursing homes,<sup>58</sup> hospitalization,<sup>59</sup> and death in the hospital<sup>59</sup> than do patients who do not use the services. Similar programs of team-coordinated home-based care exist within the VHA.<sup>60</sup>

Until comprehensive palliative care programs become widely available, the responsibility for coordination of care falls to the primary care physician, since patients and their families frequently have problems negotiating the current health care system. Table 2 details the range of palliative care services that should be considered at various stages of a serious illness; Table 3 lists services that are covered by Medicare.

---

#### AREAS OF UNCERTAINTY

---

Physicians have few data to guide them in improving their communication with patients and in their management of pain and other symptoms. The communication guidelines presented in Figure 1, although widely taught and used by palliative care experts, have not been shown to be associated with

**Table 2. Coordination of Care for Early, Middle, and Late Stages of Serious Chronic Illnesses.\***

Palliative Care Services	Early Stage	Middle Stage	Late Stage
Goals of care	Discuss diagnosis, prognosis, likely course of the illness, and disease-modifying therapies; talk about patient-centered goals, hopes, and expectations for medical treatments.	Review patient's understanding of prognosis; review efficacy and benefit-to-burden ratio for disease-modifying treatments; reassess goals of care and expectations; prepare patient and patient's family for a shift in goals; encourage paying attention to important tasks, relationships, and financial affairs.	Assess patient's understanding of diagnosis, disease course, and prognosis; review appropriateness of disease-modifying treatments; review goals of care and recommend appropriate shifts; help patient explicitly plan for a peaceful death; encourage completion of important tasks and increased attention to relationships and financial affairs.
Programmatic support	Advise patient to sign up for visiting nurse and home care services and case-management services (if available).	Advise patient to sign up for visiting nurse and home care services; consider palliative care program in hospital or at home, hospice, subacute rehabilitation, case-management services, and PACE.	Advise patient to sign up for a palliative care program in hospital or at home, case-management services, hospice, or PACE; consider nursing home placement with hospice or palliative care if patient's home caregivers are overwhelmed.
Financial planning	Advise patient to seek help in planning for financial, long-term care, and insurance needs and to begin transfer of assets if patient is considering a future Medicaid application; refer patient to a lawyer who is experienced in health issues.	Advise patient to reassess adequacy of planning for financial, medical, home care, prescription, long-term care, and family-support needs; consider hospice referral and Medicaid eligibility.	Advise patient to review all financial resources and needs; inform patient and family about financial options for personal and long-term care (e.g., hospice and Medicaid) if resources are inadequate to meet needs; explicitly recommend hospice and review its advantages; consider Medicaid eligibility.
Family support	Inform patient and family about support groups; ask about practical support needs (e.g., transportation, prescription-drug coverage, respite care, and personal care); listen to concerns.	Encourage support or counseling for family caregivers; ensure that caregivers have information about practical resources, stress, depression, and adequacy of medical care; identify respite and practical support resources; recommend help from family and friends; raise the possibility of hospice and discuss its benefits; listen to concerns.	Encourage out-of-town family to visit; refer caregivers to disease-specific support groups or counseling; inquire routinely about health, well-being, and practical needs of caregivers; offer resources for respite care; after death, send bereavement card and call after one to two weeks; screen for complicated bereavement; maintain occasional contact after patient's death; listen to concerns.

\* Early stage refers to the stage of disease at the time of diagnosis, middle stage to progressive disease and increasing functional decline, and late stage to the stage when death is imminent. PACE denotes Program of All-Inclusive Care for the Elderly.

improved outcomes. Similarly, although data are available to guide the treatment of pain due to cancer, data are lacking regarding the treatment of other pain syndromes, symptoms other than pain (such as dyspnea, fatigue, delirium, and anxiety), spiritual distress, and complicated grief and bereavement.

#### GUIDELINES

The National Comprehensive Cancer Network<sup>61</sup> has developed guidelines for patients who have advanced incurable cancer, and the National Consensus Project for Quality Palliative Care,<sup>62</sup> a collaborative effort of five national palliative care organizations, has similar guidelines for patients who

have advanced chronic illnesses. These guidelines encompass many of the recommendations detailed in this article, and the National Consensus Project guidelines also include recommendations regarding the spiritual, religious, and existential aspects of care; cultural aspects of care; and care of patients whose death is imminent.<sup>62</sup> Although the guidelines of the National Comprehensive Cancer Network are targeted to patients with terminal cancer, many patients who have early stages of the disease or uncertain prognoses or who are undergoing active curative or life-prolonging therapies can benefit from the organization's recommendations. The application of those recommendations should not be restricted to patients with a limited life expectancy.

**Table 3. Summary of Medicare Coverage for Common Services Required by Patients with Serious and Chronic Illnesses.**

Medicare Part A*		
Service	Coverage	Out-of-Pocket Cost
Ambulance	Transportation only to a hospital or skilled nursing facility and only if transportation in any other type of vehicle endangers health	20% of Medicare-approved amounts
Custodial care	No coverage for custodial care if this is the only care needed	100%
Home health care	Skilled nursing care in the home for treatment of an acute illness if at least one of the following is needed: intermittent skilled nursing care, physical or speech therapy, or continuing occupational therapy and if the patient is unable to leave the house except with major effort	None; if the patient has only Part B coverage, services are covered by Part B
Hospice	Eligibility: life expectancy of 6 mo or less if disease follows its usual course and willingness to relinquish Medicare-reimbursed services focused on prolongation of life or cure; coverage includes physician and nursing services, durable medical equipment and supplies, medications for pain or symptom control, home health aide and homemaker services (4 hr/day on average), physical and occupational therapy, short-stay hospitalizations and respite care, social work services, and bereavement services; regular Medicare covers services not related to hospice diagnosis; room and board are covered only for respite care and short hospital stays, not for nursing home or residential hospice stays	None for regular hospice services; copayment of up to \$5 for outpatient prescription drugs; 5% of Medicare-approved amount for inpatient respite care
Hospital inpatient care	General hospital services with semiprivate room	For each benefit period, deductible of \$812 for days 1–60, \$203/day for days 61–90, and \$406/day for days 91–150; benefit period begins on hospital day 1 and ends when hospital or skilled nursing care has not been received for 60 consecutive days
Skilled nursing facility or rehabilitation	If patient has Part A benefit days remaining after hospitalization; qualified hospital stay (inpatient stay of 3 consecutive days and entry into a skilled nursing facility within 30 days of discharge); need for a skilled service to treat a medical condition that was treated in the hospital or started while getting Medicare-covered skilled nursing care; no coverage for nursing home custodial care	For each benefit period, none for days 1–20, \$101.50/day for days 21–100, and 100%/day for day 101
Medicare Part B†		
Service	Coverage	Out-of-Pocket Cost
Custodial care	No coverage for custodial care if this is the only care needed	100%
Durable medical equipment	“Air-fluidized” beds, canes, commode chairs, crutches, home oxygen and equipment (with qualifications), hospital beds, nebulizers, patient lifts, suction pumps, walkers, and wheelchairs	Variable after \$100 deductible
Office visits to doctors	No coverage for routine physical and gynecologic examinations	20% of Medicare-approved rates
Outpatient mental health services	Office visits to physicians, clinical social workers, psychologists, nurse specialists, and physician assistants	50% of Medicare-approved rates
Nonphysician health care services	Office visits to clinical social workers, psychologists, physician assistants, and nurse practitioners (for medically necessary services)	20% of Medicare-approved rates
Prescription drugs	Limited coverage for erythropoietin only in end-stage renal disease, injectable drugs, limited oral cancer medications, limited oral antiemetics (if a Medicare-covered cancer drug), some infusion therapies (if considered reasonable and necessary). A Medicare discount card is currently available for other prescription drugs and a \$600 credit is available for qualifying low-income individuals. In 2005, a Medicare prescription-drug plan will become available.	In 2004, 75 to 90 percent for most prescriptions. In 2005, a \$35 monthly premium and a \$250 deductible; 25 percent of pharmacy costs between \$250 and \$2,250 in drug spending, 100 percent of costs between \$2,250 and \$3,600, and 5 percent of costs over \$3,600
Transportation	No coverage for transportation to physicians' offices, laboratory tests, outpatient physical therapy, or ambulette (van) services	100%

\* Under Part A, the patient pays no premium if the patient or the patient's spouse has paid Medicare taxes while working for 40 or more quarters.

† Under Part B, the patient pays for optional coverage with a premium of \$58.70 per month.

---

CONCLUSIONS  
AND RECOMMENDATIONS

---

The aim of palliative care is to relieve suffering and improve the quality of life for patients with advanced illnesses and their families. It is based on an interdisciplinary approach that is offered simultaneously with other appropriate medical treatments and involves close attention to the emotional, spiritual, and practical needs and goals of patients and of the people who are close to them. The patient who is described in the vignette, who has multiple chronic and serious medical conditions but is not actively dying, is ideally suited to receive high-quality palliative care. Specifically, we suggest the following approach. The patient's symptoms should be assessed and treated, as discussed in Table 1. If available, consultation with a palliative care team should be strongly considered to help with evaluating and managing his symptoms. Once the patient is comfortable, a discussion about realistic goals should be held, with particular attention given to clarifying the patient's opinion about an acceptable quality of life, identifying conditions under which the patient would consider life not worth living, establishing attainable short- and long-term goals, and designating a health care proxy. We recommend that this discussion be summarized in a treatment directive and that a health-care-proxy form be completed. Once goals are clarified, appropriate treatment and discharge planning can begin.

In the case described in the vignette, for example, the treatment goal might be to focus on the patient's comfort and on caring for him at home, and the appropriate treatments might be spoon-feeding rather than placement of a PEG tube, diuretics and

bisphosphonates, avoidance of hospitalization, and a trial of physical and occupational therapy with the aim of regaining function. The patient's discharge services will depend on the goals, the patient's insurance coverage and financial resources, and available home care services. Since a hospice program will support these goals, a referral to hospice should be considered. If hospice is not an option (e.g., because of limited home care hours or uncertainty about the prognosis), the patient should be referred to a case-management program, PACE, or a certified home care agency. The family's ability to afford medications should be evaluated, and alternative sources (such as mail-order drug plans or state-sponsored prescription plans for low-income elderly people) or less costly medications should be considered. Before discharge, a home-safety and home-needs evaluation should be performed either by an occupational therapist or through a structured interview with the patient's caregiver. An assessment of support requirements for the caregiver should be conducted to ensure that the personal care needs of the patient and the patient's family are met. Referral to a social worker can help identify community alternatives and determine Medicaid eligibility. Finally, a regular system of communication should be established between the treating physician and the home care team.

Dr. Morrison is a Paul Beeson Faculty Scholar in Aging Research, an Open Society Institute Project on Death in America Faculty Scholar, and the recipient of a Mid-Career Investigator Award in Patient-Oriented Research (K24AG022345) from the National Institute on Aging. Dr. Meier is an Open Society Institute Project on Death in America Faculty Scholar, the recipient of an Academic Career Leadership Award (K07AG00903) from the National Institute on Aging, and the director of the Center to Advance Palliative Care, a program office of the Robert Wood Johnson Foundation.

---

REFERENCES

- Field MJ, Cassel CK, eds. *Approaching death: improving care at the end of life*. Washington, D.C.: National Academy Press, 1997.
- The SUPPORT Principal Investigators. A controlled trial to improve care for seriously ill hospitalized patients: the Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments (SUPPORT). *JAMA* 1995;274:1591-8. [Erratum, *JAMA* 1996;275:1232.]
- Emanuel EJ, Fairclough DL, Slutsman J, Emanuel LL. Understanding economic and other burdens of terminal illness: the experience of patients and their caregivers. *Ann Intern Med* 2000;132:451-9.
- Emanuel EJ, Fairclough DL, Slutsman J, Alpert H, Baldwin D, Emanuel LL. Assistance from family members, friends, paid care givers, and volunteers in the care of terminally ill patients. *N Engl J Med* 1999;341:956-63.
- Covinsky KE, Goldman L, Cook EF, et al. The impact of serious illness on patients' families. *JAMA* 1994;272:1839-44.
- Christakis NA, Escarce JJ. Survival of Medicare patients after enrollment in hospice programs. *N Engl J Med* 1996;335:172-8.
- Fisher ES, Wennberg DE, Stukel TA, Gottlieb DJ, Lucas FL, Pinder EL. The implications of regional variations in Medicare spending. 2. Health outcomes and satisfaction with care. *Ann Intern Med* 2003;138:288-98.
- Sullivan AM, Lakoma MD, Block SD. The status of medical education in end-of-life care: a national report. *J Gen Intern Med* 2003;18:685-95.
- Steinhauser KE, Christakis NA, Clipp EC, McNeilly M, McIntyre L, Tulsky JA. Factors considered important at the end of life by patients, family, physicians, and other care providers. *JAMA* 2000;284:2476-82.
- Tulsky JA. Doctor-patient communication. In: Morrison RS, Meier DE, eds. *Geriatric palliative care*. New York: Oxford University Press, 2003:314-31.
- Dowsett SM, Saul JL, Butow PN, et al. Communication styles in the cancer consultation: preferences for a patient-centered approach. *Psychooncology* 2000;9:147-56.
- The EPEC Project: Education on Palliative and End-of-life Care. (Accessed April 21, 2004, at <http://www.epec.net>.)

13. Buckman R. How to break bad news: a guide for health care professionals. Baltimore: Johns Hopkins University Press, 1992.
14. Goold SD, Williams B, Arnold RM. Conflicts regarding decisions to limit treatment: a differential diagnosis. *JAMA* 2000;283:909-14.
15. Meier DE, Back AL, Morrison RS. The inner life of physicians and care of the seriously ill. *JAMA* 2001;286:3007-14.
16. Quill TE. Perspectives on care at the close of life: initiating end-of-life discussions with seriously ill patients: addressing the "elephant in the room." *JAMA* 2000;284:2502-7.
17. Fallowfield L, Jenkins V, Farewell V, Saul J, Duffy A, Eves R. Efficacy of a Cancer Research UK communication skills training model for oncologists: a randomised controlled trial. *Lancet* 2002;359:650-6.
18. Harvard Medical School Center for Palliative Care. Practical aspects in palliative medicine: integrating palliative care into clinical practice. (Accessed April 21, 2004, at <http://www.hms.harvard.edu/cdi/pallcare>.)
19. American Academy on Physician and Patient. Upcoming courses and meetings. (Accessed April 21, 2004, at <http://www.physicianpatient.org/default.htm>.)
20. Singer PA, Martin DK, Kelner M. Quality end-of-life care: patients' perspectives. *JAMA* 1999;281:163-8.
21. Finucane TE, Christmas C, Travis K. Tube feeding in patients with advanced dementia: a review of the evidence. *JAMA* 1999;282:1365-70.
22. Miles SH, Koepp R, Weber EP. Advance end-of-life treatment planning: a research review. *Arch Intern Med* 1996;156:1062-8.
23. Hammes BJ, Rooney BL. Death and end-of-life planning in one midwestern community. *Arch Intern Med* 1998;158:383-90.
24. Morrison RS, Meier DE. High rates of advance care planning in New York City's elderly. *Arch Intern Med* (in press).
25. Teno J, Lynn J, Wenger N, et al. Advance directives for seriously ill hospitalized patients: effectiveness with the Patient Self-Determination Act and the SUPPORT intervention. *J Am Geriatr Soc* 1997;45:500-7.
26. Fried TR, Bradley EH, Towle VR, Allore H. Understanding the treatment preferences of seriously ill patients. *N Engl J Med* 2002;346:1061-6.
27. Meier DE, Morrison RS. Autonomy reconsidered. *N Engl J Med* 2002;346:1087-9.
28. Doyle D, Hanks GW, Cherny NI, Calman K. Oxford textbook of palliative medicine. 3rd ed. Oxford, England: Oxford University Press, 2004.
29. TIME: Toolkit of Instruments to Measure End of life care. (Accessed April 21, 2004, at <http://www.chcr.brown.edu/pcoc/toolkit.htm>.)
30. Morrison RS, Meier DE. Geriatric palliative care. New York: Oxford University Press, 2003.
31. Bernabei R, Gambassi G, Lapane K, et al. Management of pain in elderly patients with cancer. *JAMA* 1998;279:1877-82. [Erratum, *JAMA* 1999;281:136.]
32. Bookbinder M, Coyle N, Kiss M, et al. Implementing national standards for cancer pain management: program model and evaluation. *J Pain Symptom Manage* 1996;12:334-47.
33. Manfredi PL, Morrison RS, Morris J, Goldhirsch SL, Carter JM, Meier DE. Palliative care consultations: how do they impact the care of hospitalized patients? *J Pain Symptom Manage* 2000;20:166-73.
34. Morrison RS, Magaziner J, McLaughlin MA, et al. The impact of post-operative pain on outcomes following hip fracture. *Pain* 2003;103:303-11.
35. Foley KM. Advances in cancer pain. *Arch Neurol* 1999;56:413-7.
36. AGS Panel on Persistent Pain in Older Persons. The management of persistent pain in older persons. *J Am Geriatr Soc* 2002;50:Suppl:S205-S224.
37. Luce JM, Luce JA. Perspectives on care at the close of life: management of dyspnea in patients with far-advanced lung disease: "once I lose it, it's kind of hard to catch it. . . ." *JAMA* 2001;285:1331-7.
38. Casarett DJ, Inouye SK. Diagnosis and management of delirium near the end of life. *Ann Intern Med* 2001;135:32-40.
39. Strasser F, Bruera ED. Update on anorexia and cachexia. *Hematol Oncol Clin North Am* 2002;16:589-617.
40. Block SD. Assessing and managing depression in the terminally ill patient. *Ann Intern Med* 2000;132:209-18.
41. Regnard C, Comiskey M. Nausea and vomiting in advanced cancer — a flow diagram. *Palliat Med* 1992;6:146-51.
42. McClain CS, Rosenfeld B, Breitbart W. Effect of spiritual well-being on end-of-life despair in terminally-ill cancer patients. *Lancet* 2003;361:1603-7.
43. Stroebe MS, Stroebe W. The mortality of bereavement: a review. In: Stroebe MS, Stroebe W, Hansson RO, eds. *Handbook of bereavement: theory, research, and intervention*. Cambridge, England: Cambridge University Press, 1993:175-95.
44. Ehman JW, Ott BB, Short TH, Ciampa RC, Hansen-Flaschen J. Do patients want physicians to inquire about their spiritual or religious beliefs if they become gravely ill? *Arch Intern Med* 1999;159:1803-6.
45. Mittelman MS, Ferris SH, Shulman E, Steinberg G, Levin B. A family intervention to delay nursing home placement of patients with Alzheimer disease: a randomized controlled trial. *JAMA* 1996;276:1725-31.
46. Mittelman MS, Ferris SH, Shulman E, et al. A comprehensive support program: effect on depression in spouse-caregivers of AD patients. *Gerontologist* 1995;35:792-802.
47. Stroebe MS, Stroebe W, Hansson RO. *Handbook of bereavement: theory, research, and intervention*. Cambridge, England: Cambridge University Press, 1993.
48. Christakis NA, Iwashyna TJ. The health impact of health care on families: a matched cohort study of hospice use by decedents and mortality outcomes in surviving, widowed spouses. *Soc Sci Med* 2003;57:465-75.
49. Teno JM, Clarridge BR, Casey V, et al. Family perspectives on end-of-life care at the last place of care. *JAMA* 2004;291:88-93.
50. Miller SC, Mor V, Teno J. Hospice enrollment and pain assessment and management in nursing homes. *J Pain Symptom Manage* 2003;26:791-9.
51. Center to Advance Palliative Care home page. (Accessed April 21, 2004, at <http://www.capc.org>.)
52. Finlay IG, Higginson IJ, Goodwin DM, et al. Palliative care in hospital, hospice, at home: results from a systematic review. *Ann Oncol* 2002;13:Suppl 4:257-64.
53. Smith TJ, Coyne P, Cassel B, Penberthy L, Hopson A, Hager MA. A high-volume specialist palliative care unit and team may reduce in-hospital end-of-life care costs. *J Palliat Med* 2003;6:699-705.
54. West JA, Miller NH, Parker KM, et al. A comprehensive management system for heart failure improves clinical outcomes and reduces medical resource utilization. *Am J Cardiol* 1997;79:58-63.
55. Wasson J, Gaudette C, Whaley F, Sauvigne A, Baribeau P, Welch HG. Telephone care as a substitute for routine clinic follow-up. *JAMA* 1992;267:1788-93.
56. Elkan R, Kendrick D, Dewey M, et al. Effectiveness of home based support for older people: systematic review and meta-analysis. *BMJ* 2001;323:719-25.
57. Mukamel DB, Bajorska A, Temkin-Greener H. Health care services utilization at the end of life in a managed care program integrating acute and long-term care. *Med Care* 2002;40:1136-48.
58. Eng C, Pedulla J, Eleazer GP, McCann R, Fox N. Program of All-Inclusive Care for the Elderly (PACE): an innovative model of integrated geriatric care and financing. *J Am Geriatr Soc* 1997;45:223-32.
59. Wieland D, Lamb VL, Sutton SR, et al. Hospitalization in the Program of All-Inclusive Care for the Elderly (PACE): rates, concomitants, and predictors. *J Am Geriatr Soc* 2001;49:835.]
60. Hughes SL, Weaver FM, Giobbie-Hurder A, et al. Effectiveness of team-managed home-based primary care: a randomized multicenter trial. *JAMA* 2000;284:2877-85.
61. The National Comprehensive Cancer Network. Guidelines for supportive care. (Accessed April 21, 2004, at [http://www.nccn.org/physician\\_gls/f\\_guidelines.html](http://www.nccn.org/physician_gls/f_guidelines.html).)
62. The National Consensus Project for Quality Palliative Care (NCP) home page. (Accessed April 21, 2004, at <http://www.nationalconsensusproject.org>.)

Copyright © 2004 Massachusetts Medical Society.